

Coexistence of Kochen-Specker inequalities and noncontextuality inequalities

Xiao-Dong Yu and D. M. Tong*

Department of Physics, Shandong University, Jinan 250100, China

(Dated: February 24, 2014)

Two types of inequalities, Kochen-Specker inequalities and noncontextuality inequalities, are both used to demonstrate the incompatibility between the noncontextual hidden variable model and quantum mechanics. It has been thought that noncontextuality inequalities are much more potent than Kochen-Specker inequalities, since the latter are constrained by the Kochen-Specker rules, which are regarded as an extra constraint imposed on the noncontextual hidden variable model. However, we find that a noncontextuality inequality exists in a ray set if and only if a Kochen-Specker inequality exists in the same ray set. This provides an effect approach both for constructing noncontextuality inequalities in a Kochen-Specker set and for converting a Kochen-Specker inequality to a noncontextuality inequality in any ray set.

PACS numbers: 03.65.Ta, 03.65.Ud

The noncontextual hidden variable (NCHV) model [1], as with the local hidden variable (LHV) model [2], has attracted a lot of attention. The NCHV model consists of two basic assumptions: that every observable A has a definite value $v(A)$ at all times and that $v(A)$ does not depend on whether A is measured alone or together with B or C if A is compatible with B and C . The first assumption is at the core of the hidden variable theory while the latter is the exhibition of noncontextuality. The most common observables used in the NCHV model are rays, P_i , $i = 1, 2, \dots$. The NCHV model indicates that $v(P_i)$ can take only 0 or 1.

To examine the inconsistency of the NCHV model with quantum mechanics, the pioneering works [1, 3–8] used to impose the constraint of preserving the algebraic structure of compatible observables, i.e., the so-called product rule $v(AB) = v(A)v(B)$ and sum rule $v(A + B) = v(A) + v(B)$ for two compatible observables A and B . For rays, the constraint is equivalent to the Kochen-Specker (KS) rules: $v(P_i)v(P_j) = 0$ if P_i and P_j are orthogonal, and exactly one $v(P_i)$ is 1 out of an orthogonal basis set of rays. A ray set $\mathcal{S}$ is called a KS set if all possible value assignments to the ray set violate the KS rules. By employing the KS rules, Kochen and Specker [1] illustrated logically that the NCHV model is inconsistent with quantum mechanics, which is usually called the KS theorem. Their original argument involved a KS set of 117 rays in a three-dimensional Hilbert space. The proof of the KS theorem is simplified step by step [3–7], with smaller KS sets, e.g., with 31 rays in a three-dimensional space by Conway and Kochen [5] or with 18 rays in a four-dimensional space by Cabello *et al.* [7]. To simplify further the logical proof, KS inequalities, as a reformulation of the KS theorem, were then proposed. Hereafter, when we use the term KS inequalities, we refer to those inequalities in which the value assignments are constrained both by the two basic assumptions in the

NCHV model and by the KS rules [9]. The statistical approach proposed by Clifton [8] may be regarded as the embryonic form of a KS inequality. Klyachko *et al.* [10] constructed a KS inequality by using a set of only five rays in a three-dimensional space. More KS inequalities can be found in recent papers [9, 11–16].

Since KS inequalities are not only based on the assumption of noncontextuality, but are also dependent on the KS rules, which is an extra assumption imposed on the NCHV model and difficult to test in experiment, researchers prefer to find inequalities without employing the KS rules. Such general inequalities, in which the value assignments are constrained only by the two basic assumptions in the NCHV model but not by the KS rules, are called noncontextuality inequalities in order to differentiate them from KS inequalities [9]. Klyachko *et al.* [10] found one such inequality, the KCBS inequality. Cabello [17] constructed several state-independent noncontextuality inequalities, which are violated by all quantum states. Soon afterwards, Badziąg *et al.* [18] put forward a general approach to construct a state-independent noncontextuality inequality from a KS set, which shows that any KS set can always lead to a noncontextuality inequality. Yu and Oh [14] found a 13-ray state-independent noncontextuality inequality in a non-KS set. Some of these theoretical results have been demonstrated in experiments [19–29].

As discussed above, both KS inequalities and noncontextuality inequalities have been used to demonstrate the incompatibility between the NCHV model and quantum mechanics. It has been usually thought that noncontextuality inequalities are much more potent than KS inequalities, since the KS rules are regarded as an extra constraint imposed on the NCHV model. However, it is much easier to construct a KS inequality than a noncontextuality inequality in a ray set, since a great number of the value assignments to the set can be removed by the KS rules. Interestingly, we notice that in some cases such as those in Refs. [9, 10, 14], both a KS inequality and a noncontextuality inequality exist in the same ray set. The result in Ref. [18] also implies that noncontext-

* tdm@sdu.edu.cn

tuality inequalities and KS inequalities coexist in a KS set. This started us wondering whether a noncontextuality inequality always coexists with a KS inequality in any ray set, not only in some particular ray sets, and whether the former can be derived by the aid of the latter if the coexistence is true. In this Rapid Communication, we address these issues. We will prove the following theorem.

Theorem: A noncontextuality inequality exists in a ray set if and only if a Kochen-Specker inequality exists in the same ray set.

The theorem uncovers an essential relationship between KS inequalities and noncontextuality inequalities. It clarifies that a noncontextuality inequality always coexists with a KS inequality in any ray set. The proof of theorem provides an effective approach both for constructing a noncontextuality inequality in a KS set and for converting a KS inequality to a noncontextuality inequality in any ray set.

We now prove the theorem. To this end, we will first show that a noncontextuality inequality can be derived from a given KS inequality.

Consider an n -dimensional quantum system. Let $\mathcal{S}$ be a μ -ray set of the system, $\mathcal{S} = \{P_1, P_2, \dots, P_\mu\}$. P_i have definite values 0 or 1 in the NCHV model, while they should be viewed as rank-1 projective operators $\hat{P}_i$ in quantum mechanics. Suppose that a KS inequality has been given as

$$\langle F \rangle_{ks} \leq f, \quad (1)$$

where $F \equiv F(P_1, P_2, \dots, P_\mu)$ is a function of the observables $P_1, P_2, \dots, P_\mu$, $\langle F \rangle_{ks}$ represents the average value of the function in the NCHV model, and f is a constant number. Hereafter, the subscript ks in $\langle \dots \rangle_{ks}$ means that $\langle \dots \rangle_{ks}$ is calculated under the constraint of the KS rules, i.e., the value assignments to the ray set $\{P_1, P_2, \dots, P_\mu\}$ obey the KS rules.

With the above knowledge, we may now start to derive a noncontextuality inequality from inequality (1).

First, we introduce a new function, $\tilde{F} = \tilde{F}(P_1, P_2, \dots, P_\mu)$. The ray set $\mathcal{S}$ can be illustrated by a graph $\mathcal{G}$. Each vertex of $\mathcal{G}$ corresponds to a ray of $\mathcal{S}$, and two vertices are linked by an edge if and only if the corresponding rays are orthogonal. Besides, we use $\mathcal{S}^\alpha$, $\alpha = 1, 2, \dots, L$, to denote the α th subset of the rays that can form an n -dimensional orthogonal basis, $\mathcal{S}^\alpha = \{P_{k_1^\alpha}, P_{k_2^\alpha}, \dots, P_{k_n^\alpha}\}$, where $k_1^\alpha < k_2^\alpha < \dots < k_n^\alpha$. L is the total number of such subsets found in $\mathcal{S}$, and it is zero if $\mathcal{S}$ does not contain any n -dimensional orthogonal basis. The new function $\tilde{F}$ is defined as

$$\tilde{F} = \lambda F - \sum_{\substack{i>j=1, \\ \hat{P}_i \hat{P}_j = 0}}^{\mu} P_i P_j + \sum_{\alpha=1}^L \left(\sum_{i=1}^n P_{k_i^\alpha} - 2 \sum_{i>j=1}^n P_{k_i^\alpha} P_{k_j^\alpha} \right), \quad (2)$$

where λ is a positive parameter to be determined. The first term in $\tilde{F}$ is the function F defined in Eq. (1), the

second term corresponds to the pairs of rays that are orthogonal, and the third term corresponds to the sets of rays that form an n -dimensional orthogonal basis.

Second, we calculate the average value of $\tilde{F}$ under the NCHV model, with

$$\begin{aligned} \langle \tilde{F} \rangle = & \lambda \langle F \rangle - \sum_{\substack{i>j=1, \\ \hat{P}_i \hat{P}_j = 0}}^{\mu} \langle P_i P_j \rangle \\ & + \sum_{\alpha=1}^L \left(\sum_{i=1}^n \langle P_{k_i^\alpha} \rangle - 2 \sum_{i>j=1}^n \langle P_{k_i^\alpha} P_{k_j^\alpha} \rangle \right). \end{aligned} \quad (3)$$

Clearly, $\langle \tilde{F} \rangle$ depends on the values of P_i and $P_i P_j$. We first consider the case where the value assignments to the ray set obey the KS rules. Under the KS rules, $P_i P_j = 0$ if P_i and P_j are orthogonal, and exactly one $P_i = 1$ out of an orthogonal basis. We immediately have

$$\tilde{F}|_{ks} = \lambda F|_{ks} + L, \quad (4)$$

where $\tilde{F}|_{ks}$ ($F|_{ks}$) is the value of function $\tilde{F}$ (F). The subscript ks means that $\tilde{F}|_{ks}$ ($F|_{ks}$) is obtained when the assignments of values 0, 1 to the ray set obey the KS rules. Noting that Eq. (1) implies $F|_{ks} \leq f$, we further have

$$\tilde{F}|_{ks} \leq \lambda f + L. \quad (5)$$

We now consider the case where the value assignments to the ray set violate the KS rules. The value of the expression $(\sum_{i=1}^n P_{k_i^\alpha} - 2 \sum_{i>j=1}^n P_{k_i^\alpha} P_{k_j^\alpha})$, denoted by $(\dots)_\alpha$, is only dependent on the number of 1's (or 0's) assigned to the rays in the subset $\mathcal{S}^\alpha$ but independent of particular value assignments due to the symmetry in the expression. Thus $(\dots)_\alpha = x(2-x)$, if the number of 1's in the value assignments is x . It shows that $(\dots)_\alpha \leq 0$ if $x \neq 1$. Therefore, if the KS rules are violated, at least one $P_i P_j = 1$ or one $(\dots)_\alpha \leq 0$. We then have

$$\tilde{F}|_{exks} \leq \lambda F|_{exks} + L - 1, \quad (6)$$

where the subscript $exks$ means that the values are obtained in the case where the value assignments to the ray set $\mathcal{S}$ violate the KS rules. Let $f' = \max\{F|_{exks}\}$. We further have

$$\tilde{F}|_{exks} \leq \lambda f' + L - 1. \quad (7)$$

Inequality (5) is valid for the value assignments obeying the KS rules, and inequality (7) is valid for the value assignments violating the KS rules. By combining Eqs. (5) and (7), we finally obtain the inequality

$$\tilde{F} \leq \max\{\lambda f + L, \lambda f' + L - 1\}. \quad (8)$$

Equation (8) is always valid for all possible value assignments to the ray set, irrespective of whether the KS rules are obeyed or violated. Hence, we get an inequality without assuming the KS rules,

$$\langle \tilde{F} \rangle \leq \max\{\lambda f + L, \lambda f' + L - 1\}. \quad (9)$$

Third, we discuss the quantum violation of the inequality. In quantum mechanics, each observable P_i corresponds to an operator $\hat{P}_i = |\varphi_i\rangle\langle\varphi_i|$. $\langle P_i P_j \rangle = 0$ if P_i and P_j are orthogonal, and $\sum_{i=1}^n \langle P_i \rangle = \langle \sum_{i=1}^n P_i \rangle = 1$ if $\{P_i, i = 1, 2, \dots, n\}$ form an orthogonal basis. In quantum mechanics, the average value of $\tilde{F}$, expressed as Eq. (3), reads

$$\langle \tilde{F} \rangle_\psi = \lambda \langle F \rangle_\psi + L, \quad (10)$$

where the subscript ψ indicates that the averages $\langle \tilde{F} \rangle_\psi$ and $\langle F \rangle_\psi$ are calculated by using the theory of quantum mechanics for state $|\psi\rangle$. By comparing Eqs. (9) and (10), quantum violation occurs if

$$\lambda \langle F \rangle_\psi + L > \max\{\lambda f + L, \lambda f' + L - 1\}. \quad (11)$$

The validity of Eq. (11) relies on the parameters f , f' , and λ . If both $f \geq f'$ and $\lambda > 0$ or if both $f < f'$ and $0 < \lambda \leq \frac{1}{f'-f}$, Eq. (11) can be guaranteed by

$$\langle F \rangle_\psi > f, \quad (12)$$

which is just the quantum violation of Eq. (1). Therefore, there always exists λ such that the quantum violation of Eq. (1) can always guarantee the quantum violation of Eq. (9).

So far, we have proved that a KS inequality can always lead to a noncontextuality inequality, and the latter is violated in quantum mechanics if the former is violated. Note that the value assignments of variables in a KS inequality are constrained by both the assumption of noncontextuality and the KS rules while the assignments in a noncontextuality inequality are constrained only by the assumption of noncontextuality. The value assignments satisfying both the assumption of noncontextuality and the KS rules are a subset of the value assignments satisfying only the assumption of noncontextuality. If an inequality is valid for all the value assignments satisfying the assumption of noncontextuality, it must be valid for the subset of the value assignments, too. Therefore, a noncontextuality inequality itself can be regarded as a KS inequality. This completes the proof of the theorem that a noncontextuality inequality exists if and only if a KS inequality exists in a ray set.

It is interesting to see that the above proof has actually provided an effective approach for constructing a noncontextuality inequality with the aid of a KS inequality in any ray set. It is generally easier to construct a KS inequality than a noncontextuality inequality in a ray set. Once a KS inequality is constructed, the theorem can convert it to a noncontextuality inequality. By following the proof, all the well-known KS inequalities proposed in the literature can be converted to noncontextuality inequalities. For example, we consider the first inequality proposed by Yu and Oh [14], with a 13-ray set in a three-dimensional quantum system. The state-independent KS inequality [see Eq. (2) in Ref. [14]] reads $\langle F \rangle_{ks} \leq 1$, with $F = \sum_{i=1}^4 P_i$, where we have rewritten the expressions using

our notations. Here, $P_1, P_2, \dots, P_{13}$ are corresponding to $h_0, h_2, h_2, h_3, z_2, y_2^+, y_2^-, z_3, y_3^+, y_3^-, z_1, y_1^+, y_1^-$ in Ref. [14], respectively. Starting from it, we can immediately obtain a state-independent noncontextuality inequality by following the process in the proof of the theorem. Indeed, by comparing it with Eq. (1), we have $f = 1$, and by using the definition $f' = \max\{F|_{exks}\}$, we have $f' = 4$, which indicate that $0 < \lambda \leq \frac{1}{f'-f} = \frac{1}{3}$. The state-independent noncontextuality inequality is given by Eq. (9) as $\langle \tilde{F} \rangle \leq 4 + \lambda$, where $\tilde{F}$ is defined as Eq. (2) with $n = 3$, $\mu = 13$, $L = 4$ [30], $0 < \lambda \leq \frac{1}{3}$, and $F = \sum_{i=1}^4 P_i$.

As shown in the above example, one may easily convert a KS inequality to a noncontextuality inequality, expressed as Eq. (9), by calculating f' and λ . However, the bound in expression (9) may not be tight, since it has been relaxed for the generality during the proof of the theorem. In a specific application, the tight bound, $\max\{\tilde{F}_{ks}, \tilde{F}_{exks}\}$, can be obtained by detailedly calculating $\tilde{F}_{ks}$ and $\tilde{F}_{exks}$, and a tight inequality is given as $\langle \tilde{F} \rangle \leq \max\{\tilde{F}_{ks}, \tilde{F}_{exks}\}$, instead of Eq. (9). We would like to illustrate this point by considering the KS inequality proposed by Klyachko *et al.* in Ref. [10], with a five-ray set in a three-dimensional quantum system. The state-dependent KS inequality [see Eq. (7) in Ref. [10]] reads $\langle F \rangle_{ks} \leq 2$, with $F = \sum_{i=1}^5 P_i$. Since the last term in Eq. (2) vanishes due to $L = 0$ for this instance, Eq. (2) is reduced to $\tilde{F} = \lambda F - \sum_{i=1}^5 P_i P_{i+1}$, where the subscript i is modulo 5. In this case, we have $f = 2$, $f' = \max\{F|_{exks}\} = 5$, and $0 < \lambda \leq \frac{1}{f'-f} = \frac{1}{3}$. Substituting them into Eq. (9), we obtain $\langle \tilde{F} \rangle \leq \max\{2\lambda, 5\lambda - 1\}$ with $0 < \lambda \leq \frac{1}{3}$, which is a state-dependent noncontextuality inequality but not a tight one. To find the tight bound, we calculate $\tilde{F}_{ks}$ and $\tilde{F}_{exks}$ detailedly, and have $\tilde{F}|_{ks} = 0, \lambda, 2\lambda, \tilde{F}|_{exks} = 2\lambda - 1, 3\lambda - 1, 3\lambda - 2, 4\lambda - 3, 5\lambda - 5$, and $\max\{\tilde{F}\} = \max\{\tilde{F}|_{ks}, \tilde{F}|_{exks}\} = \max\{2\lambda, 3\lambda - 1, 4\lambda - 3, 5\lambda - 5\}$. We then obtain the tight noncontextuality inequality, $\langle \tilde{F} \rangle \leq \max\{2\lambda, 3\lambda - 1, 4\lambda - 3, 5\lambda - 5\}$ with $0 < \lambda \leq 1$, where $0 < \lambda \leq 1$ is determined by requiring the quantum violation of the inequality. For $\lambda = 1$, the inequality reads $\sum_{i=1}^5 \langle P_i \rangle - \sum_{i=1}^5 \langle P_i P_{i+1} \rangle \leq 2$, which is exactly the KCBS inequality given in Ref. [10] if P_i is replaced by A_i with $P_i = \frac{1}{2}(1 - A_i)$.

It should be noted that the above discussion is concerned essentially with a general non-KS set. If the ray set under consideration is a KS set, then the process of obtaining a noncontextuality inequality becomes much simpler. By definition, for a KS set, all possible value assignments to the set violate the KS rules. In this case, all value assignments obeying the KS rules certainly satisfy any given inequalities logically, because there are no value assignments obeying the KS rules at all, and therefore Eq. (1) can be any form of inequality with a quantum violation. This implies that a state-independent noncontextuality inequality always exists for a KS set. However, it also implies that Eq. (1) is useless to a KS set due to the arbitrariness of its form. Fortunately, the proof of the

theorem provides a simple approach to obtain a state-independent noncontextuality inequality from a KS set. In fact, for a KS set, $\tilde{F}$ can be directly defined as

$$\tilde{F} = - \sum_{\substack{i>j=1, \\ \tilde{P}_i \tilde{P}_j=0}}^{\mu} P_i P_j + \sum_{\alpha=1}^L \left(\sum_{i=1}^n P_{k_i^\alpha} - 2 \sum_{i>j=1}^n P_{k_i^\alpha} P_{k_j^\alpha} \right), \quad (13)$$

and there is

$$\tilde{F}|_{\text{exks}} \leq L - 1. \quad (14)$$

Noting that expression (5) does not need to be considered since all the value assignments to a KS set violate the KS rules, we then have

$$\langle \tilde{F} \rangle \leq L - 1. \quad (15)$$

Equation (15) is a state-independent noncontextuality inequality, since $\langle \tilde{F} \rangle_\psi = L$ ($> L - 1$) for any quantum state $|\psi\rangle$. Comparing with the previous method for constructing a noncontextuality inequality from a KS set, proposed by Badziąg *et al.* [18], our approach is much simpler. In the previous method, one needs to enlarge the given KS set to construct a state-independent noncontextuality inequality in general and correlations of n compatible observables are involved, while only correlations of two compatible observables are involved in the present approach.

In passing, we point out that P_i can be replaced by $A_i = 1 - 2P_i$ in the proof of the theorem, and all of the above discussions work for A_i although we have

used P_i , instead of A_i , in this Rapid Communication for simplicity. By substituting $P_i = \frac{1}{2}(1 - A_i)$ and $P_i P_j = \frac{1}{4}(1 - A_i - A_j + A_i A_j)$, $\tilde{F}$ in Eqs. (2) and (13) can be written as a function of A_i , and the same procedure can be followed. However, the expressions as well as the calculations are much simpler by using P_i than A_i . Besides, although our theoretical approach can help to determine the existence of a KS inequality in a ray set, the inequality obtained may not be optimal. The numerical approach proposed in Ref. [31] may be used to optimize it if needed.

In conclusion, we have established a theorem that a noncontextuality inequality exists if and only if a KS inequality exists in a ray set. The theorem not only solves the coexistence of the two types of inequalities in any ray set, but also provides an effective approach both for constructing noncontextuality inequalities in a KS set and for converting a KS inequality to a noncontextuality inequality in any ray set. A noncontextuality inequality is immediately obtained when the approach is applied to a KS set, and all the known KS inequalities can be converted to noncontextuality inequalities when it is applied to any ray set.

ACKNOWLEDGMENTS

We thank Murray Batchelor for useful discussions. This work was supported by NSF China by Grant No.11175105 and the Taishan Scholarship Project of Shandong Province. D.M.T acknowledges support from the National Basic Research Program of China by Grant No. 2009CB929400.

-
- [1] S. Kochen and E. Specker, J. Math. Mech. **17**, 59 (1967).
 - [2] J. S. Bell, Rev. Mod. Phys. **38**, 447 (1966).
 - [3] A. Peres, J. Phys. A **24**, L175 (1991).
 - [4] J. Bub, Found. Phys. **26**, 787 (1996).
 - [5] J. H. Conway and J. Kochen, First reported in A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer, Dordrecht, 1993), p. 114. See also J. H. Conway and S. Kochen, in *Quantum [Un]speakeables: From Bell to Quantum Information*, edited by R. A. Bertlmann and A. Zeilinger (Springer, Berlin, 2002), p. 257.
 - [6] M. Kernaghan, J. Phys. A **27**, L829 (1994).
 - [7] A. Cabello, J. Estebarez, and G. García-Alcaine, Phys. Lett. A **212**, 183 (1996).
 - [8] R. Clifton, Am. J. Phys. **61**, 443 (1993).
 - [9] I. Bengtsson, K. Blanchfield, and A. Cabello, Phys. Lett. A **376**, 374 (2012).
 - [10] A. A. Klyachko, M. A. Can, S. Binicioğlu, and A. S. Shumovsky, Phys. Rev. Lett. **101**, 020403 (2008).
 - [11] C. Simon, Č. Brukner, and A. Zeilinger, Phys. Rev. Lett. **86**, 4427 (2001).
 - [12] J.-Å. Larsson, Europhys. Lett. **58**, 799 (2002).
 - [13] A. Cabello, S. Filipp, H. Rauch, and Y. Hasegawa, Phys. Rev. Lett. **100**, 130404 (2008).
 - [14] S. Yu and C. H. Oh, Phys. Rev. Lett. **108**, 030402 (2012).
 - [15] A. Cabello, S. Severini, and A. Winter, arXiv:1010.2163.
 - [16] P. Kurzyński and D. Kaszlikowski, Phys. Rev. A **86**, 042125 (2012).
 - [17] A. Cabello, Phys. Rev. Lett. **101**, 210401 (2008).
 - [18] P. Badziąg, I. Bengtsson, A. Cabello, and I. Pitowsky, Phys. Rev. Lett. **103**, 050401 (2009).
 - [19] M. Michler, H. Weinfurter, and M. Żukowski, Phys. Rev. Lett. **84**, 5457 (2000).
 - [20] Y.-F. Huang, C.-F. Li, Y.-S. Zhang, J.-W. Pan, and G.-C. Guo, Phys. Rev. Lett. **90**, 250401 (2003).
 - [21] Y. Hasegawa, R. Loidl, G. Badurek, M. Baron, and H. Rauch, Phys. Rev. Lett. **97**, 230401 (2006).
 - [22] H. Bartosik, J. Klepp, C. Schmitzer, S. Sponar, A. Cabello, H. Rauch, and Y. Hasegawa, Phys. Rev. Lett. **103**, 040403 (2009).
 - [23] B. H. Liu, Y. F. Huang, Y. X. Gong, F. W. Sun, Y. S. Zhang, C. F. Li, and G. C. Guo, Phys. Rev. A **80**, 044101 (2009).
 - [24] G. Kirchmair, F. Zähringer, R. Gerritsma, M. Kleinmann, O. Gühne, A. Cabello, R. Blatt, and C. Roos, Nature (London) **460**, 494 (2009).
 - [25] E. Amsalem, M. Rådmark, M. Bourennane, and A. Cabello, Phys. Rev. Lett. **103**, 160405 (2009).
 - [26] O. Moussa, C. A. Ryan, D. G. Cory, and R. Laflamme, Phys. Rev. Lett. **104**, 160501 (2010).

- [27] R. Lapkiewicz, P. Li, C. Schaeff, N. Langford, S. Ramelow, M. Wieśniak, and A. Zeilinger, *Nature (London)* **474**, 490 (2011).
- [28] C. Zu, Y.-X. Wang, D.-L. Deng, X.-Y. Chang, K. Liu, P.-Y. Hou, H.-X. Yang, and L.-M. Duan, *Phys. Rev. Lett.* **109**, 150401 (2012).
- [29] X. Zhang, M. Um, J. Zhang, S. An, Y. Wang, D.-l. Deng, C. Shen, L.-M. Duan, and K. Kim, *Phys. Rev. Lett.* **110**, 070401 (2013).
- [30] There are four subsets of the rays that form a three-dimensional orthogonal basis, $\mathcal{S}^1 = \{P_5, P_6, P_7\}$, $\mathcal{S}^2 = \{P_8, P_9, P_{10}\}$, $\mathcal{S}^3 = \{P_{11}, P_{12}, P_{13}\}$, and $\mathcal{S}^4 = \{P_5, P_8, P_{11}\}$.
- [31] M. Kleinmann, C. Budroni, J.-A. Larsson, O. Gühne, and A. Cabello, *Phys. Rev. Lett.* **109**, 250402 (2012).